

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 5/00, A61K 38/00	A2	(11) International Publication Number: WO 99/25727 (43) International Publication Date: 27 May 1999 (27.05.99)
(21) International Application Number: PCT/US98/24210 (22) International Filing Date: 13 November 1998 (13.11.98) (30) Priority Data: 60/065,442 14 November 1997 (14.11.97) US (71) Applicant (for all designated States except US): AMYLIN PHARMACEUTICALS, INC. [US/US]; 9373 Towne Centre Drive, San Diego, CA 92121 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BEELEY, Nigel, Robert, Arnold [US/US]; 227 Loma Corta Drive, Solana Beach, CA 92131 (US). PRICKETT, Kathryn, S. [US/US]; 7612 Trailbrush Terrace, San Diego, CA 92126 (US). (74) Agents: CONSALVI, Mary, S. et al.; Lyon & Lyon LLP, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With declaration under Article 17(2)(a); without abstract; title not checked by the International Searching Authority.</i>
(54) Title: NOVEL EXENDIN AGONIST COMPOUNDS		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America ..
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DESCRIPTION

NOVEL EXENDIN AGONIST COMPOUNDS

Related Applications

This application claims the benefit of U.S. Provisional Application No. 60/065,442, filed November 14, 1997, the contents of which are hereby incorporated by reference in their entirety.

Field of the Invention

The present invention relates to novel compounds which have activity as exendin agonists. These compounds are useful in treatment of Type I and II diabetes, in treatment of disorders which would be benefited by agents which lower plasma glucose levels and in treatment of disorders which would be benefited with agents useful in delaying and/or slowing gastric emptying.

BACKGROUND

The following description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art to the presently claimed invention, nor that any of the publications specifically or implicitly referenced are prior art to that invention.

Exendin

The exendins are peptides that are found in venom of the Gila-monster, a lizard endogenous to Arizona and Northern Mexico. Exendin-3 [SEQ. ID. NO. 1] is present in the venom of *Heloderma horridum*, and exendin-4 [SEQ. ID. NO. 2] is present in the venom of *Heloderma suspectum* (Eng, J., et al.,

J. Biol. Chem., 265:20259-62, 1990; Eng., J., et al., J. Biol. Chem., 267:7402-05, 1992). The amino acid sequence of exendin-3 is shown in Figure 1. The amino acid sequence of exendin-4 is shown in Figure 2. The exendins have some sequence similarity to several members of the glucagon-like peptide family, with the highest homology, 53%, being to GLP-1[7-36]NH₂ [SEQ. ID. NO. 3] (Goke, et al., J. Biol. Chem., 268:19650-55, 1993). GLP-1[7-36]NH₂, also known as proglucagon[78-107] or simply "GLP-1" as used most often herein, has an insulintropic effect, stimulating insulin secretion from pancreatic β -cells; GLP-1 also inhibits glucagon secretion from pancreatic α -cells (Ørsov, et al., Diabetes, 42:658-61, 1993; D'Alessio, et al., J. Clin. Invest., 97:133-38, 1996). The amino acid sequence of GLP-1 is shown in Figure 3. GLP-1 is reported to inhibit gastric emptying (Willms B, et al., J Clin Endocrinol Metab 81 (1): 327-32, 1996; Wettergren A, et al., Dig Dis Sci 38 (4): 665-73, 1993), and gastric acid secretion. Schjoldager BT, et al., Dig Dis Sci 34 (5): 703-8, 1989; O'Halloran DJ, et al., J Endocrinol 126 (1): 169-73, 1990; Wettergren A, et al., Dig Dis Sci 38 (4): 665-73, 1993). GLP-1[7-37], which has an additional glycine residue at its carboxy terminus, also stimulates insulin secretion in humans (Ørsov, et al., Diabetes, 42:658-61, 1993).

A transmembrane G-protein adenylate-cyclase-coupled receptor believed to be responsible for the insulintropic effect of GLP-1 has been cloned from a β -cell line (Thorens, Proc. Natl. Acad. Sci. USA 89:8641-45, 1992), herein after referred to as the "cloned GLP-1 receptor." Exendin-4 is reportedly acts at GLP-1 receptors on insulin-secreting β TC1 cells, at dispersed acinar cells from guinea pig pancreas, and at parietal cells from stomach; the peptide is also

reported to stimulate somatostatin release and inhibit gastrin release in isolated stomachs (Goke, et al., J. Biol. Chem. 268:19650-55, 1993; Schepp, et al., Eur. J. Pharmacol., 69:183-91, 1994; Eissele, et al., Life Sci., 55:629-34, 1994). Exendin-3 and exendin-4 were reportedly found to stimulate cAMP production in, and amylase release from, pancreatic acinar cells (Malhotra, R., et al., Regulatory Peptides, 41:149-56, 1992; Raufman, et al., J. Biol. Chem. 267:21432-37, 1992; Singh, et al., Regul. Pept. 53:47-59, 1994). Based on their insulinotropic activities, the use of exendin-3 and exendin-4 for the treatment of diabetes mellitus and the prevention of hyperglycemia has been proposed (Eng, U.S. Patent No. 5,424,286).

Agents which serve to delay gastric emptying have found a place in medicine as diagnostic aids in gastro-intestinal radiologic examinations. For example, glucagon is a polypeptide hormone which is produced by the α cells of the pancreatic islets of Langerhans. It is a hyperglycemic agent which mobilizes glucose by activating hepatic glycogenolysis.

It can to a lesser extent stimulate the secretion of pancreatic insulin. Glucagon is used in the treatment of insulin-induced hypoglycemia, for example, when administration of glucose intravenously is not possible. However, as glucagon reduces the motility of the gastro-intestinal tract it is also used as a diagnostic aid in gastro-intestinal radiological examinations. Glucagon has also been used in several studies to treat various painful gastro-intestinal disorders associated with spasm. Daniel, et al. (Br. Med. J., 3:720, 1974) reported quicker symptomatic relief of acute diverticulitis in patients treated with glucagon compared with those who had been treated with analgesics or antispasmodics. A review by Glauser, et al. (J. Am. Coll. Emergency Physns, 8:228, 1979)

described relief of acute esophageal food obstruction following glucagon therapy. In another study glucagon significantly relieved pain and tenderness in 21 patients with biliary tract disease compared with 22 patients treated with placebo (M.J. Stower, et al., Br. J. Surg., 69:591-2, 1982).

Methods for regulating gastrointestinal motility using amylin agonists are described in International Application No. PCT/US94/10225, published March 16, 1995.

Methods for regulating gastrointestinal motility using exendin agonists are described in U.S. Patent Application Serial No. 08/908,867, filed August 8, 1997 entitled "Methods for Regulating Gastrointestinal Motility," which application is a continuation-in-part of U.S. Patent Application Serial No. 08/694,954 filed August 8, 1996.

Methods for reducing food intake using exendin agonists are described in U.S. Patent Application Serial No. 09/003,869, filed January 7, 1998, entitled "Use of Exendin and Agonists Thereof for the Reduction of Food Intake," which claims the benefit of U.S. Provisional Application Nos. 60/034,905, filed January 7, 1997, 60/055,404, filed August 7, 1997, 60/065,442, filed November 14, 1997 and 60/066,029, filed November 14, 1997.

Novel exendin agonist compounds are described in PCT Application Serial No. PCT/US98/16387 filed August 6, 1998, entitled "Novel Exendin Agonist Compounds," claiming the benefit of U.S. Provisional Application Serial No. 60/055,404, filed August 8, 1997. Other novel exendin agonist compounds are described in PCT Application Serial No. _____ filed November 13, 1998, entitled "Novel Exendin Agonist Compounds," which claims the benefit of U.S. Provisional Application No. 60/066,029 filed November 14, 1997.

SUMMARY OF THE INVENTION

According to one aspect, the present invention provides novel exendin agonist compounds which exhibit advantageous properties which include effects in slowing gastric emptying and lowering plasma glucose levels.

According to the present invention, provided are compounds of the formula (I) [SEQ. ID. NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁; wherein

Xaa₁ is His, Arg or Tyr;

Xaa₂ is Ser, Gly, Ala or Thr;

Xaa₃ is Asp or Glu;

Xaa₅ is Ala or Thr;

Xaa₆ is Ala, Phe, Tyr or naphthylalanine;

Xaa₇ is Thr or Ser;

Xaa₈ is Ala, Ser or Thr;

Xaa₉ is Asp or Glu;

Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;

Xaa₁₁ is Ala or Ser;

Xaa₁₂ is Ala or Lys;

Xaa₁₃ is Ala or Gln;

Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;

Xaa₁₅ is Ala or Glu;

Xaa₁₆ is Ala or Glu;

Xaa₁₇ is Ala or Glu;

Xaa₁₉ is Ala or Val;

Xaa₂₀ is Ala or Arg;

Xaa₂₁ is Ala or Leu;

Xaa₂₂ is Phe, Tyr or naphthylalanine;
Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
or Met;
Xaa₂₄ is Ala, Glu or Asp;
Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
Xaa₂₆ is Ala or Leu;
Xaa₂₇ is Ala or Lys;
Xaa₂₈ is Ala or Asn;
Z₁ is -OH,

-NH₂

Gly-Z₂,

Gly Gly-Z₂,

Gly Gly Xaa₃₁-Z₂,

Gly Gly Xaa₃₁ Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂;

Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently Pro,

homoproline, 3Hyp, 4Hyp, thioproline,

N-alkylglycine, N-alkylpentylglycine or

N-alkylalanine; and

Z₂ is -OH or -NH₂;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈,
Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀,
Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈ are Ala. Also
included within the scope of the present invention are
pharmaceutically acceptable salts of the compounds of formula
(I) and pharmaceutical compositions including said compounds
and salts thereof.

Also within the scope of the present invention are narrower genera of compounds having peptides of various lengths, for example genera of compounds which do not include peptides having a length of 28, 29 or 30 amino acid residues, respectively. Additionally, the present invention includes narrower genera of compounds having particular amino acid sequences, for example, compounds of the formula (I) [SEQ. ID. NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₈ Xaa₁₉
Xaa₂₀ Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁;

wherein

Xaa₁ is His or Arg;

Xaa₂ is Gly or Ala;

Xaa₃ is Asp or Glu;

Xaa₅ is Ala or Thr;

Xaa₆ is Ala, Phe or naphthylalanine;

Xaa₇ is Thr or Ser;

Xaa₈ is Ala, Ser or Thr;

Xaa₉ is Asp or Glu;

Xaa₁₀ is Ala, Leu or pentylglycine;

Xaa₁₁ is Ala or Ser;

Xaa₁₂ is Ala or Lys;

Xaa₁₃ is Ala or Gln;

Xaa₁₄ is Ala, Leu or pentylglycine;

Xaa₁₅ is Ala or Glu;

Xaa₁₆ is Ala or Glu;


Xaa₁₇ is Ala or Glu;

Xaa₁₉ is Ala or Val;

Xaa₂₀ is Ala or Arg;

Xaa₂₁ is Ala or Leu;
 Xaa₂₂ is Phe or naphthylalanine;
 Xaa₂₃ is Ile, Val or tert-butylglycine;
 Xaa₂₄ is Ala, Glu or Asp;
 Xaa₂₅ is Ala, Trp, or Phe;
 Xaa₂₆ is Ala or Leu;
 Xaa₂₇ is Ala or Lys;
 Xaa₂₈ is Ala or Asn;
 Z₁ is -OH,
 -NH₂,
 Gly-Z₂,
 Gly Gly -Z₂,
 Gly Gly Xaa₃₁-Z₂,
 Gly Gly Xaa₃₁ Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or Gly Gly
 Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂;
 Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected
 from the group consisting of Pro, homoproline,
 thioprolin and N-methylalanine; and
 Z₂ is -OH or -NH₂;
 provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈,
 Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀,
 Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈ are Ala; and
 pharmaceutically acceptable salts thereof.

Also provided are compounds of the formula (II) [SEQ.
 ID. NO. 66]:



Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ X₁ -Z₁; wherein

Xaa₁ is His, Arg, Tyr or 4-imidazopropionyl;

Xaa₂ is Ser, Gly, Ala or Thr;

Xaa₃ is Asp or Glu;

Xaa₅ is Ala or Thr;

Xaa₆ is Ala, Phe, Tyr or naphthylalanine;

Xaa₇ is Thr or Ser;

Xaa₈ is Ala, Ser or Thr;

Xaa₉ is Asp or Glu;

Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;

Xaa₁₁ is Ala or Ser;

Xaa₁₂ is Ala or Lys;

Xaa₁₃ is Ala or Gln;

Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;

Xaa₁₅ is Ala or Glu;

Xaa₁₆ is Ala or Glu;

Xaa₁₇ is Ala or Glu;

Xaa₁₉ is Ala or Val;

Xaa₂₀ is Ala or Arg;

Xaa₂₁ is Ala, Leu or Lys-NH^t-R where R is Lys, Arg, C₁-C₁₀
straight chain or branched alkanoyl or cycloalkylalkanoyl;

Xaa₂₂ is Phe, Tyr or naphthylalanine;

Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
or Met;

Xaa₂₄ is Ala, Glu or Asp;

Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;

Xaa₂₆ is Ala or Leu;

X₁ is Lys Asn, Asn Lys, Lys-NH^t-R Asn, Asn Lys-NH^t-R, Lys-NH^t-R

Ala, Ala Lys-NH^t-R where R is Lys, Arg, C₁-C₁₀ straight chain
or branched alkanoyl or cycloalkylalkanoyl

Z_1 is -OH,
 -NH₂,
 Gly-Z₂,
 Gly Gly-Z₂,
 Gly Gly Xaa₃₁-Z₂,
 Gly Gly Xaa₃₁ Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂;

wherein

Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently
 selected from the group consisting of Pro,
 homoproline, 3Hyp, 4Hyp, thioproline,
 N-alkylglycine, N-alkylpentylglycine and
 N-alkylalanine; and

Z_2 is -OH or -NH₂;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈,
 Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀,
 Xaa₂₁, Xaa₂₄, Xaa₂₅, and Xaa₂₆ are Ala. Also within the scope of
 the present invention are pharmaceutically acceptable salts
 of the compounds of formula (II) and pharmaceutical
 compositions including said compounds and salts thereof.

Preferred exendin agonist compounds of formula (II)
 include those wherein Xaa₁ is His, Tyr or 4-imidazopropionyl.

More preferably Xaa₁ is His. Also, preferred are those
 compounds of formula (II) wherein Xaa₁ is 4-imidazopropionyl.

Preferred are those compounds of formula (II) wherein
 Xaa₂ is Gly.

Preferred compounds of formula (II) are those wherein
 Xaa₁₄ is Leu, pentylglycine or Met.

Preferred compounds of formula (II) are those wherein Xaa₂₅ is Trp or Phe.

According to one aspect, preferred are compounds of formula (II) wherein Xaa₆ is Phe or naphthylalanine; Xaa₂₂ is Phe or naphthylalanine; and Xaa₂₃ is Ile or Val. More preferably, Z₁ is -NH₂. According to one aspect, especially preferred are such compounds of formula (II) wherein Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected from the group consisting of Pro, homoproline, thioproline and N-alkylalanine. More preferably, Z₂ is -NH₂.

Preferred compounds of formula (II) include those wherein X₁ is Lys Asn, Lys-NH^f-R Asn, or Lys-NH^f-R Ala where R is Lys, Arg, C₁-C₁₀ straight chain or branched alkanoyl. Especially preferred compounds of formula (II) include Compound Nos. 62-69 [SEQ. ID. NOS. 67-74].

Definitions


In accordance with the present invention and as used herein, the following terms are defined to have the following meanings, unless explicitly stated otherwise.

The term "amino acid" refers to natural amino acids, unnatural amino acids, and amino acid analogs, all in their D and L stereoisomers if their structure allow such stereoisomeric forms. Natural amino acids include alanine (Ala), arginine (Arg), asparagine (Asn), aspartic acid (Asp), cysteine (Cys), glutamine (Gln), glutamic acid (Glu), glycine (Gly), histidine (His), isoleucine (Ile), leucine (Leu), Lysine (Lys), methionine (Met), phenylalanine (Phe), proline (Pro), serine (Ser), threonine (Thr), typtophan (Trp), tyrosine (Tyr) and valine (Val). Unnatural amino acids include, but are not limited to azetidinecarboxylic acid, 2-aminoadipic acid, 3-aminoadipic acid, beta-alanine, aminopropionic acid, 2-aminobutyric acid, 4-aminobutyric

acid, 6-aminocaproic acid, 2-aminoheptanoic acid, 2-aminoisobutyric acid, 3-aminoisobutyric acid, 2-aminopimelic acid, tertiary-butylglycine, 2,4-diaminoisobutyric acid, desmosine, 2,2'-diaminopimelic acid, 2,3-diaminopropionic acid, N-ethylglycine, N-ethylasparagine, homoproline, hydroxylysine, allo-hydroxylysine, 3-hydroxyproline, 4-hydroxyproline, isodesmosine, allo-isoleucine, N-methylalanine, N-methylglycine, N-methylisoleucine, N-methylpentylglycine, N-methylvaline, naphthalanine, norvaline, norleucine, ornithine, pentylglycine, pipecolic acid and thioproline. Amino acid analogs include the natural and unnatural amino acids which are chemically blocked, reversibly or irreversibly, or modified on their N-terminal amino group or their side-chain groups, as for example, methionine sulfoxide, methionine sulfone, S-(carboxymethyl)-cysteine, S-(carboxymethyl)-cysteine sulfoxide and S-(carboxymethyl)-cysteine sulfone.

The term "amino acid analog" refers to an amino acid wherein either the C-terminal carboxy group, the N-terminal amino group or side-chain functional group has been chemically codified to another functional group. For example, aspartic acid-(beta-methyl ester) is an amino acid analog of aspartic acid; N-ethylglycine is an amino acid analog of glycine; or alanine carboxamide is an amino acid analog of alanine.

The term "amino acid residue" refers to radicals having the structure: (1) $-C(O)-R-NH-$, wherein R typically is $-CH(R')-$, wherein R' is an amino acid side chain, typically H or a carbon containing substituent; or (2)



wherein p is 1, 2 or 3 representing the azetidinecarboxylic acid, proline or pipecolic acid residues, respectively.

The term "lower" referred to herein in connection with organic radicals such as alkyl groups defines such groups with up to and including about 6, preferably up to and including 4 and advantageously one or two carbon atoms. Such groups may be straight chain or branched chain.

"Pharmaceutically acceptable salt" includes salts of the compounds of the present invention derived from the combination of such compounds and an organic or inorganic acid. In practice the use of the salt form amounts to use of the base form. The compounds of the present invention are useful in both free base and salt form, with both forms being considered as being within the scope of the present invention.

In addition, the following abbreviations stand for the following:

- "ACN" or "CH₃CN" refers to acetonitrile.
- "Boc", "tBoc" or "Tboc" refers to t-butoxy carbonyl.
- "DCC" refers to N,N'-dicyclohexylcarbodiimide.
- "Fmoc" refers to fluorenylmethoxycarbonyl.
- "HBTU" refers to 2-(1H-benzotriazol-1-yl)-1,1,3,3,-tetramethyluronium hexafluorophosphate.
- "HOBt" refers to 1-hydroxybenzotriazole monohydrate.
- "homop" or hPro" refers to homoproline.
- "MeAla" or "Nme" refers to N-methylalanine.
- "naph" refers to naphthylalanine.

"pG" or pGly" refers to pentylglycine.

"tBuG" refers to tertiary-butylglycine.

"ThioP" or tPro" refers to thioproline.

"3Hyp" refers to 3-hydroxyproline

"4Hyp" refers to 4-hydroxyproline

"NAG" refers to N-alkylglycine

"NAPG" refers to N-alkylpentylglycine

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the amino acid sequence for exendin-3 [SEQ. ID. NO. 1].

Figure 2 depicts the amino acid sequence for exendin-4 [SEQ. ID. NO. 2].

Figure 3 depicts the amino acid sequence for GLP-1[7-36]NH₂ (GLP-1) [SEQ. ID. NO. 3].

Figure 4 depicts the amino acid sequences for certain compounds of the present invention, Compounds 1-61 [SEQ. ID. NOS. 5 to 65].

Figure 5 depicts the effect on lowering blood glucose of various concentrations of Compound 3 [SEQ. ID. NO. 7].

Figure 6 depicts the effect on gastric emptying of various concentrations of Compound 2 [SEQ. ID. NO. 6].

Figure 7 depicts a comparison of effects on gastric emptying of various concentrations of Compound 3 [SEQ. ID. NO. 7].

Figure 8 depicts a comparison of effects on gastric emptying of various concentrations of Compound 10 [SEQ. ID. NO. 14].

Figure 9 depicts a comparison of effects on gastric emptying of various concentrations of Compound 13 [SEQ. ID. NO. 17].

Figure 10 depicts the amino acid sequences for certain compounds of the present invention, Compound Nos. 62 - 69 [SEQ. ID. NOS. 67 - 74].

DETAILED DESCRIPTION OF THE INVENTION

According to the present invention, provided are compounds of the formula (I) [SEQ. ID. NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁; wherein
Xaa₁ is His, Arg or Tyr;
Xaa₂ is Ser, Gly, Ala or Thr;
Xaa₃ is Asp or Glu;
Xaa₅ is Ala or Thr;
Xaa₆ is Ala, Phe, Tyr or naphthylalanine;
Xaa₇ is Thr or Ser;
Xaa₈ is Ala, Ser or Thr;
Xaa₉ is Asp or Glu;
Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
Xaa₁₁ is Ala or Ser;
Xaa₁₂ is Ala or Lys;
Xaa₁₃ is Ala or Gln;
Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
Xaa₁₅ is Ala or Glu;
Xaa₁₆ is Ala or Glu;
Xaa₁₇ is Ala or Glu;
Xaa₁₉ is Ala or Val;
Xaa₂₀ is Ala or Arg;
Xaa₂₁ is Ala or Leu;
Xaa₂₂ is Phe, Tyr or naphthylalanine;

Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
or Met;

Xaa₂₄ is Ala, Glu or Asp;

Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;

Xaa₂₆ is Ala or Leu;

Xaa₂₇ is Ala or Lys;

Xaa₂₈ is Ala or Asn;

Z₁ is -OH,

-NH₂

Gly-Z₂,

Gly Gly-Z₂,

Gly Gly Xaa₃₁-Z₂,

Gly Gly Xaa₃₁ Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂;

Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently Pro,
homoproline, 3Hyp, 4Hyp, thioproline,

N-alkylglycine, N-alkylpentylglycine or

N-alkylalanine; and

Z₂ is -OH or -NH₂;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈,
Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀,
Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈ are Ala. Also within
the scope of the present invention are pharmaceutically
acceptable salts of the compounds for formula (I) and
pharmaceutical compositions including said compounds and
salts thereof.

Preferred N-alkyl groups for N-alkylglycine, N-
alkylpentylglycine and N-alkylalanine include lower alkyl

groups preferably of 1 to about 6 carbon atoms, more preferably of 1 to 4 carbon atoms. Suitable compounds of formula (I) include those identified in Examples 1-61 ("Compounds 1-61," respectively) [SEQ. ID. NOS. 5 to 65], as well as those corresponding compounds identified in Examples 70 and 71.

Preferred such exendin agonist compounds include those wherein Xaa₁ is His or Tyr. More preferably Xaa₁ is His.

Preferred are those such compounds wherein Xaa₂ is Gly.

Preferred are those such compounds wherein Xaa₁₄ is Leu, pentylglycine or Met.

Preferred such compounds are those wherein Xaa₂₅ is Trp or Phe.

Preferred compounds of formula (I) are those where Xaa₆ is Phe or naphthylalanine; Xaa₂₂ is Phe or naphthylalanine and Xaa₂₃ is Ile or Val.

Preferred are compounds of formula (I) wherein Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected from Pro, homoproline, thioproline and N-alkylalanine.

Preferably Z₁ is -NH₂.

Preferable Z₂ is -NH₂.

According to one aspect, preferred are compounds of formula (I) wherein Xaa₁ is His or Tyr, more preferably His; Xaa₂ is Gly; Xaa₆ is Phe or naphthylalanine; Xaa₁₄ is Leu, pentylglycine or Met; Xaa₂₂ is Phe or naphthylalanine; Xaa₂₃ is Ile or Val; Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected from Pro, homoproline, thioproline or N-alkylalanine. More preferably Z₁ is -NH₂.

According to an especially preferred aspect, especially preferred compounds include those of formula (I) wherein: Xaa₁ is His or Arg; Xaa₂ is Gly or Ala; Xaa₃ is Asp or Glu; Xaa₅ is Ala or Thr; Xaa₆ is Ala, Phe or naphthylalanine; Xaa₇ is Thr or Ser; Xaa₈ is Ala, Ser or Thr; Xaa₉ is Asp or Glu;

Xaa₁₀ is Ala, Leu or pentylglycine; Xaa₁₁ is Ala or Ser; Xaa₁₂ is Ala or Lys; Xaa₁₃ is Ala or Gln; Xaa₁₄ is Ala, Leu or pentylglycine; Xaa₁₅ is Ala or Glu; Xaa₁₆ is Ala or Glu; Xaa₁₇ is Ala or Glu; Xaa₁₉ is Ala or Val; Xaa₂₀ is Ala or Arg; Xaa₂₁ is Ala or Leu; Xaa₂₂ is Phe or naphthylalanine; Xaa₂₃ is Ile, Val or tert-butylglycine; Xaa₂₄ is Ala, Glu or Asp; Xaa₂₅ is Ala, Trp or Phe; Xaa₂₆ is Ala or Leu; Xaa₂₇ is Ala or Lys; Xaa₂₈ is Ala or Asn; Z₁ is -OH, -NH₂, Gly-Z₂, Gly Gly-Z₂, Gly Gly Xaa₃₁-Z₂, Gly Gly Xaa₃₁ Ser-Z₂, Gly Gly Xaa₃₁ Ser Ser-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂, Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂; Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ being independently Pro homoproline, thioproline or N-methylalanine; and Z₂ being -OH or -NH₂; provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈ are Ala. Especially preferred compounds of formula (I) include those having the amino acid sequence of SEQ. ID. NOS. 6-27 (Compounds 2-23).

According to an especially preferred aspect, provided are compounds of compound (I) where Xaa₁₄ is Leu, Ile, Val or pentylglycine, more preferably Leu or pentylglycine, and Xaa₂₅ is Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds will be less susceptible to oxidative degradation, both in vitro and in vivo, as well as during synthesis of the compound.

Also within the scope of the present invention are narrower genera of compounds having peptides of various lengths, for example genera of compounds which do not include peptides having a length of 28, 29 or 30 amino acid residues, respectively. Additionally, the present invention includes narrower genera of compounds having particular amino acid

sequences, for example, compounds of the formula (I) [SEQ. ID. NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₈ Xaa₁₉
Xaa₂₀ Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁;

wherein

Xaa₁ is His or Arg;
Xaa₂ is Gly or Ala;
Xaa₃ is Asp or Glu;
Xaa₅ is Ala or Thr;
Xaa₆ is Ala, Phe or naphthylalanine;
Xaa₇ is Thr or Ser;
Xaa₈ is Ala, Ser or Thr;
Xaa₉ is Asp or Glu;
Xaa₁₀ is Ala, Leu or pentylglycine;
Xaa₁₁ is Ala or Ser;
Xaa₁₂ is Ala or Lys;
Xaa₁₃ is Ala or Gln;
Xaa₁₄ is Ala, Leu or pentylglycine;
Xaa₁₅ is Ala or Glu;
Xaa₁₆ is Ala or Glu;
Xaa₁₇ is Ala or Glu;
Xaa₁₉ is Ala or Val;
Xaa₂₀ is Ala or Arg;
Xaa₂₁ is Ala or Leu;
Xaa₂₂ is Phe or naphthylalanine;
Xaa₂₃ is Ile, Val or tert-butylglycine;
Xaa₂₄ is Ala, Glu or Asp;
Xaa₂₅ is Ala, Trp, or Phe;
Xaa₂₆ is Ala or Leu;

Xaa₂₇ is Ala or Lys;

Xaa₂₈ is Ala or Asn;

Z₁ is -OH,

 $-\text{NH}_2,$

Gly-Z₂,

Gly Gly -Z₂,

Gly Gly Xaa₃₁-Z₂,

Gly Gly Xaa₃₁ Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or Gly Gly

Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂;

Xaa_{31} , Xaa_{36} , Xaa_{37} and Xaa_{38} are independently selected

from the group consisting of Pro, homoproline,

thioprolin and N-methylalalanine; and

Z_2 is $-OH$ or $-NH_2$;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈ are Ala; and pharmaceutically acceptable salts thereof.

Also provided are compounds of the formula (II) [SEQ. ID. NO. 66]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ X₁ -Z₁; wherein

Xaa₁ is His, Arg or Tyr or 4-imidazopropionyl;

Xaa, is Ser, Gly, Ala or Thr;

Xaa₃ is Asp or Glu;

Xaa₅ is Ala or Thr;
 Xaa₆ is Ala, Phe, Tyr or naphthylalanine;
 Xaa₇ is Thr or Ser;
 Xaa₈ is Ala, Ser or Thr;
 Xaa₉ is Asp or Glu;
 Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
 Xaa₁₁ is Ala or Ser;
 Xaa₁₂ is Ala or Lys;
 Xaa₁₃ is Ala or Gln;
 Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
 Xaa₁₅ is Ala or Glu;
 Xaa₁₆ is Ala or Glu;
 Xaa₁₇ is Ala or Glu;
 Xaa₁₉ is Ala or Val;
 Xaa₂₀ is Ala or Arg;
 Xaa₂₁ is Ala, Leu or Lys-NH^ε-R where R is Lys, Arg, C₁-C₁₀ straight chain or branched alkanoyl or cycloalkylalkanoyl;
 Xaa₂₂ is Phe, Tyr or naphthylalanine;
 Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met;
 Xaa₂₄ is Ala, Glu or Asp;
 Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
 Xaa₂₆ is Ala or Leu;
 X₁ is Lys Asn, Asn Lys, Lys-NH^ε-R Asn, Asn Lys-NH^ε-R, Lys-NH^ε-R Ala, Ala Lys-NH^ε-R where R is Lys, Arg, C₁-C₁₀ straight chain or branched alkanoyl or cycloalkylalkanoyl
 Z₁ is -OH,
 -NH₂,
 Gly-Z₂,
 Gly Gly-Z₂,
 Gly Gly Xaa₃₁-Z₂,
 Gly Gly Xaa₃₁ Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or
Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂;

wherein

Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected from the group consisting of Pro, homoproline, 3Hyp, 4Hyp, thioproline, N-alkylglycine, N-alkylpentylglycine and N-alkylalanine; and Z₂ is -OH or -NH₂;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, and Xaa₂₆ are Ala. Also within the scope of the present invention are pharmaceutically acceptable salts of the compound of formula (II) and pharmaceutical compositions including said compounds and salts thereof.

Preferred exendin agonist compounds of formula (II) include those wherein Xaa₁ is His, Tyr or 4-imidazopropionyl. More preferably Xaa₁ is His.

Preferred are those compounds of formula (II) wherein Xaa₁ is 4-imidazopropionyl.

Preferred are those compounds of formula (II) wherein Xaa₂ is Gly.

Preferred compounds of formula (II) are those wherein Xaa₁₄ is Leu, pentylglycine or Met.

Preferred compounds of formula (II) are those wherein Xaa₂₅ is Trp or Phe.

According to one aspect, preferred are compounds of formula (II) wherein Xaa₆ is Phe or naphthylalanine; and Xaa₂₂ is Phe or naphthylalanine; and Xaa₂₃ is Ile or Val. More

preferably, Z_1 is $-NH_2$. According to one aspect, especially preferred are such compounds of formula (II) wherein Xaa_{31} , Xaa_{36} , Xaa_{37} and Xaa_{38} are independently selected from the group consisting of Pro, homoproline, thioproline and N-alkylalanine. More preferably, Z_2 is $-NH_2$.

Preferred compounds of formula (II) include those wherein X_1 is Lys Asn, Lys- NH^e -R Asn, or Lys- NH^e -R Ala where R is Lys, Arg, C_1 - C_{10} straight chain or branched alkanoyl. Preferred compounds of formula (II) include Compound Nos. 62-69 [SEQ. ID. NOS. 67-74].

The compounds referenced above form salts with various inorganic and organic acids and bases. Such salts include salts prepared with organic and inorganic acids, for example, HCl, HBr, H_2SO_4 , H_3PO_4 , trifluoroacetic acid, acetic acid, formic acid, methanesulfonic acid, toluenesulfonic acid, maleic acid, fumaric acid and camphorsulfonic acid. Salts prepared with bases include ammonium salts, alkali metal salts, e.g. sodium and potassium salts, and alkali earth salts, e.g. calcium and magnesium salts. Acetate, hydrochloride, and trifluoroacetate salts are preferred. The salts may be formed by conventional means, as by reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion on a suitable ion exchange resin.

Utility

The compounds described above are useful in view of their pharmacological properties. In particular, the compounds of the invention are exendin agonists, and possess

activity as agents to regulate gastric motility and to slow gastric emptying, as evidenced by the ability to reduce post-prandial glucose levels in mammals.

The compounds of the present invention are useful in in vitro and in vivo scientific methods for investigation of exendins and exendin agonists for example in methods such as those described in Examples A-E below.

Preparation of Compounds

The compounds of the present invention may be prepared using standard solid-phase peptide synthesis techniques and preferably an automated or semiautomated peptide synthesizer.

Typically, using such techniques, an α -N-carbamoyl protected amino acid and an amino acid attached to the growing peptide chain on a resin are coupled at room temperature in an inert solvent such as dimethylformamide, N-methylpyrrolidinone or methylene chloride in the presence of coupling agents such as dicyclohexylcarbodiimide and 1-hydroxybenzotriazole in the presence of a base such as diisopropylethylamine. The α -N-carbamoyl protecting group is removed from the resulting peptide-resin using a reagent such as trifluoroacetic acid or piperidine, and the coupling reaction repeated with the next desired N-protected amino acid to be added to the peptide chain. Suitable N-protecting groups are well known in the art, with t-butyloxycarbonyl (tBoc) and fluorenylmethoxycarbonyl (Fmoc) being preferred herein.

The solvents, amino acid derivatives and 4-methylbenzhydryl-amine resin used in the peptide synthesizer may be purchased from Applied Biosystems Inc. (Foster City, CA). The following side-chain protected amino acids may be purchased from Applied Biosystems, Inc.: Boc-Arg(Mts), Fmoc-Arg(Pmc), Boc-Thr(Bzl), Fmoc-Thr(t-Bu), Boc-Ser(Bzl), Fmoc-Ser(t-Bu), Boc-Tyr(BrZ), Fmoc-Tyr(t-Bu), Boc-

Lys(Cl-Z), Fmoc-Lys(Boc), Boc-Glu(Bzl), Fmoc-Glu(t-Bu), Fmoc-His(Trt), Fmoc-Asn(Trt), and Fmoc-Gln(Trt). Boc-His(BOM) may be purchased from Applied Biosystems, Inc. or Bachem Inc. (Torrance, CA). Anisole, dimethylsulfide, phenol, ethanedithiol, and thioanisole may be obtained from Aldrich Chemical Company (Milwaukee, WI). Air Products and Chemicals (Allentown, PA) supplies HF. Ethyl ether, acetic acid and methanol may be purchased from Fisher Scientific (Pittsburgh, PA).

Solid phase peptide synthesis may be carried out with an automatic peptide synthesizer (Model 430A, Applied Biosystems Inc., Foster City, CA) using the NMP/HOBt (Option 1) system and tBoc or Fmoc chemistry (see, Applied Biosystems User's Manual for the ABI 430A Peptide Synthesizer, Version 1.3B July 1, 1988, section 6, pp. 49-70, Applied Biosystems, Inc., Foster City, CA) with capping. Boc-peptide-resins may be cleaved with HF (-5°C to 0°C, 1 hour). The peptide may be extracted from the resin with alternating water and acetic acid, and the filtrates lyophilized. The Fmoc-peptide resins may be cleaved according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc., 1990, pp. 6-12). Peptides may be also be assembled using an Advanced Chem Tech Synthesizer (Model MPS 350, Louisville, Kentucky).

Peptides may be purified by RP-HPLC (preparative and analytical) using a Waters Delta Prep 3000 system. A C4, C8 or C18 preparative column (10 μ , 2.2 x 25 cm; Vydac, Hesperia, CA) may be used to isolate peptides, and purity may be determined using a C4, C8 or C18 analytical column (5 μ , 0.46 x 25 cm; Vydac). Solvents (A=0.1% TFA/water and B=0.1% TFA/CH₃CN) may be delivered to the analytical column at a flowrate of 1.0 ml/min and to the preparative column at 15 ml/min. Amino acid analyses may be performed on the Waters

Pico Tag system and processed using the Maxima program. Peptides may be hydrolyzed by vapor-phase acid hydrolysis (115°C, 20-24 h). Hydrolysates may be derivatized and analyzed by standard methods (Cohen, et al., The Pico Tag Method: A Manual of Advanced Techniques for Amino Acid Analysis, pp. 11-52, Millipore Corporation, Milford, MA (1989)). Fast atom bombardment analysis may be carried out by M-Scan, Incorporated (West Chester, PA). Mass calibration may be performed using cesium iodide or cesium iodide/glycerol. Plasma desorption ionization analysis using time of flight detection may be carried out on an Applied Biosystems Bio-Ion 20 mass spectrometer. Electrospray mass spectroscopy may be carried and on a VG-Trio machine.

Peptide compounds useful in the invention may also be prepared using recombinant DNA techniques, using methods now known in the art. See, e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d Ed., Cold Spring Harbor (1989). Non-peptide compounds useful in the present invention may be prepared by art-known methods.

Formulation and Administration

Compounds of the invention are useful in view of their exendin-like effects, and may conveniently be provided in the form of formulations suitable for parenteral (including intravenous, intramuscular and subcutaneous) or nasal, sublingual, buccal or oral administration. In some cases, it will be convenient to provide an exendin agonist and another anti-gastric-emptying agent, such as glucagon, an amylin, or an amylin agonist, in a single composition or solution for administration together. In other cases, it may be more advantageous to administer another anti-emptying agent separately from said exendin agonist. In yet other cases, it may be beneficial to provide an exendin agonist either co-

formulated or separately with other glucose lowering agents such as insulin. A suitable administration format may best be determined by a medical practitioner for each patient individually. Suitable pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, e.g., Remington's Pharmaceutical Sciences by E.W. Martin. See also Wang, Y.J. and Hanson, M.A. "Parenteral Formulations of Proteins and Peptides: Stability and Stabilizers," Journal of Parenteral Science and Technology, Technical Report No. 10, Supp. 42:2S (1988).

Compounds useful in the invention can be provided as parenteral compositions for injection or infusion. They can, for example, be suspended in an inert oil, suitably a vegetable oil such as sesame, peanut, olive oil, or other acceptable carrier. Preferably, they are suspended in an aqueous carrier, for example, in an isotonic buffer solution at a pH of about 5.6 to 7.4. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH buffering agents. Useful buffers include for example, sodium acetate/acetic acid buffers. A form of repository or "depot" slow release preparation may be used so that therapeutically effective amounts of the preparation are delivered into the bloodstream over many hours or days following transdermal injection or other form of delivery.

The desired isotonicity may be accomplished using sodium chloride or other pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol, polyols (such as mannitol and sorbitol), or other inorganic or organic solutes. Sodium chloride is preferred particularly for buffers containing sodium ions.

The claimed compounds can also be formulated as pharmaceutically acceptable salts (e.g., acid addition salts) and/or complexes thereof. Pharmaceutically acceptable salts are non-toxic salts at the concentration at which they are administered. The preparation of such salts can facilitate the pharmacological use by altering the physical-chemical characteristics of the composition without preventing the composition from exerting its physiological effect. Examples of useful alterations in physical properties include lowering the melting point to facilitate transmucosal administration and increasing the solubility to facilitate the administration of higher concentrations of the drug.

Pharmaceutically acceptable salts include acid addition salts such as those containing sulfate, hydrochloride, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, cyclohexylsulfamate and quinate. Pharmaceutically acceptable salts can be obtained from acids such as hydrochloric acid, sulfuric acid, phosphoric acid, sulfamic acid, acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, *p*-toluenesulfonic acid, cyclohexylsulfamic acid, and quinic acid. Such salts may be prepared by, for example, reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion on a suitable ion exchange resin.

Carriers or excipients can also be used to facilitate administration of the compound. Examples of carriers and excipients include calcium carbonate, calcium phosphate,

various sugars such as lactose, glucose, or sucrose, or types of starch, cellulose derivatives, gelatin, vegetable oils, polyethylene glycols and physiologically compatible solvents.

The compositions or pharmaceutical composition can be administered by different routes including intravenously, intraperitoneal, subcutaneous, and intramuscular, orally, topically, transdermally, buccally or transmucosally.

If desired, solutions of the above compositions may be thickened with a thickening agent such as methyl cellulose. They may be prepared in emulsified form, either water in oil or oil in water. Any of a wide variety of pharmaceutically acceptable emulsifying agents may be employed including, for example, acacia powder, a non-ionic surfactant (such as a Tween), or an ionic surfactant (such as alkali polyether alcohol sulfates or sulfonates, e.g., a Triton).

Compositions useful in the invention are prepared by mixing the ingredients following generally accepted procedures. For example, the selected components may be simply mixed in a blender or other standard device to produce a concentrated mixture which may then be adjusted to the final concentration and viscosity by the addition of water or thickening agent and possibly a buffer to control pH or an additional solute to control tonicity.

For use by the physician, the compounds will be provided in dosage unit form containing an amount of an exendin agonist, with or without another anti-emptying agent. Therapeutically effective amounts of an exendin agonist for use in the control of gastric emptying and in conditions in which gastric emptying is beneficially slowed or regulated are those that decrease post-prandial blood glucose levels, preferably to no more than about 8 or 9 mM or such that blood glucose levels are reduced as desired. In diabetic or glucose intolerant individuals, plasma glucose levels are

higher than in normal individuals. In such individuals, beneficial reduction or "smoothing" of post-prandial blood glucose levels, may be obtained. As will be recognized by those in the field, an effective amount of therapeutic agent will vary with many factors including the patient's physical condition, the blood sugar level or level of inhibition of gastric emptying to be obtained, and other factors.

Such pharmaceutical compositions are useful in causing gastric hypomotility in a subject and may be used as well in other disorders where gastric motility is beneficially reduced.

The effective daily anti-emptying dose of the compounds will typically be in the range of 0.001 or 0.005 to about 5 mg/day, preferably about 0.01 or 0.05 to 2 mg/day and more preferably about 0.05 or 0.1 to 1 mg/day, for a 70 kg patient. The exact dose to be administered is determined by the attending clinician and is dependent upon where the particular compound lies within the above quoted range, as well as upon the age, weight and condition of the individual.

Administration should begin at the first sign of symptoms or shortly after diagnosis of diabetes mellitus. Administration may be by injection, preferably subcutaneous or intramuscular, or by other routes for example, by oral, nasal, sublingual or buccal administration. For oral, nasal or buccal dosages should be increased to about 5-10 fold over injection dosages.

Generally, in treating or preventing elevated, inappropriate, or undesired post-prandial blood glucose levels, the compounds of this invention may be administered to patients in need of such treatment in a dosage ranges similar to those given above, however, the compounds are administered more frequently, for example, one, two, or three times a day.

The optimal formulation and mode of administration of compounds of the present application to a patient depend on factors known in the art such as the particular disease or disorder, the desired effect, and the type of patient. While the compounds will typically be used to treat human patients, they may also be used to treat similar or identical diseases in other vertebrates such as other primates, farm animals such as swine, cattle and poultry, and sports animals and pets such as horses, dogs and cats.

To assist in understanding the present invention the following Examples are included which describe the results of a series of experiments. The experiments relating to this invention should not, of course, be construed as specifically limiting the invention and such variations of the invention, now known or later developed, which would be within the purview of one skilled in the art are considered to fall within the scope of the invention as described herein and hereinafter claimed.

EXAMPLE 1

Preparation of Compound 1

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly-
NH₂ [SEQ. ID. NO. 5]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.). In general, single-coupling cycles were used throughout the synthesis and Fast Moc (HBTU activation) chemistry was

employed. Deprotection (Fmoc group removal) of the growing peptide chain was achieved using piperidine. Final deprotection of the completed peptide resin was achieved using a mixture of triethylsilane (0.2 mL), ethanedithiol (0.2 mL), anisole (0.2 mL), water (0.2 mL) and trifluoroacetic acid (15 mL) according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc.) The peptide was precipitated in ether/water (50 mL) and centrifuged. The precipitate was reconstituted in glacial acetic acid and lyophilized. The lyophilized peptide was dissolved in water). Crude purity was about 75%.

Used in purification steps and analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). The solution containing peptide was applied to a preparative C-18 column and purified (10% to 40% Solvent B in Solvent A over 40 minutes). Purity of fractions was determined isocratically using a C-18 analytical column. Pure fractions were pooled furnishing the above-identified peptide. Analytical RP-HPLC (gradient 30% to 50% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 18.9 minutes. Electrospray Mass Spectrometry (M): calculated 3408.0; found 3408.9.

EXAMPLE 2

Preparation of Compound 2

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂ [SEQ.
ID. NO. 6]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 40% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 17.9 minutes. Electrospray Mass Spectrometry (M): calculated 3294.7; found 3294.8.

EXAMPLE 3

Preparation of Compound 3

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 7]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 29% to 36% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 20.7 minutes. Electrospray Mass Spectrometry (M): calculated 3237.6; found 3240.

EXAMPLE 4Preparation of Compound 4

His Ala Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 8]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 15.2 minutes. Electrospray Mass Spectrometry (M): calculated 3251.6; found 3251.5.

EXAMPLE 5Preparation of Compound 5

His Gly Glu Gly Ala Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 9]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved

from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 13.1 minutes. Electrospray Mass Spectrometry (M): calculated 3207.6; found 3208.3.

EXAMPLE 6

Preparation of Compound 6

His Gly Glu Gly Thr Ala Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 10]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 35% to 45% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 12.8 minutes. Electrospray Mass Spectrometry (M): calculated 3161.5; found 3163.

EXAMPLE 7Preparation of Compound 7

His Gly Glu Gly Thr Phe Thr Ala Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 11]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 15.2 minutes. Electrospray Mass Spectrometry (M): calculated 3221.6; found 3222.7.

EXAMPLE 8Preparation of Compound 8

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 12]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to

Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 34% to 44% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3199.4.

EXAMPLE 9

Preparation of Compound 9

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ala Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 13]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 15.7 minutes. Electrospray Mass Spectrometry (M): calculated 3221.6; found 3221.6.

EXAMPLE 10

Preparation of Compound 10

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Ala Gln Leu Glu⁺
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.

ID. NO. 14]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 18.1 minutes. Electrospray Mass Spectrometry (M): calculated 3180.5; found 3180.9.

EXAMPLE 11

Preparation of Compound 11

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Ala Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 15]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 17.0 minutes. Electrospray Mass Spectrometry (M): calculated 3180.6; found 3182.8.

EXAMPLE 12Preparation of Compound 12

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Ala Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 16]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.9 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3195.9.

EXAMPLE 13Preparation of Compound 13

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Ala
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 17]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved

from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 17.9 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3179.0.

EXAMPLE 14

Preparation of Compound 14

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Ala Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 18]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.0.

EXAMPLE 15Preparation of Compound 15

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Ala Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 19]

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 13.7 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3179.0.

EXAMPLE 16Preparation of Compound 16

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Ala Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 20]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to

Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 35% to 45% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.0 minutes. Electrospray Mass Spectrometry (M): calculated 3209.6; found 3212.8.

EXAMPLE 17

Preparation of Compound 17

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Ala Leu Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 21]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3152.5; found 3153.5.

EXAMPLE 18

Preparation of Compound 18

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu⁺
Glu Glu Ala Val Arg Ala Phe Ile Glu Phe Leu Lys Asn-NH₂ [SEQ.

ID. NO. 22]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 35% to 45% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 12.1 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3197.7.

EXAMPLE 19

Preparation of Compound 19

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Ala Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 23]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 10.9 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.5.

EXAMPLE 20Preparation of Compound 20

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Ala Leu Lys Asn-NH₂ [SEQ.
ID. NO. 24]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 17.5 minutes. Electrospray Mass Spectrometry (M): calculated 3161.5; found 3163.0.

EXAMPLE 21Preparation of Compound 21

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Ala Lys Asn-NH₂ [SEQ.
ID. NO. 25]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc⁻-protected amino acids (Applied Biosystems, Inc.), cleaved

from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 19.5 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3199.

EXAMPLE 22

Preparation of Compound 22

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Ala Asn-NH₂ [SEQ.
ID. NO. 26]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.5 minutes. Electrospray Mass Spectrometry (M): calculated 3180.5; found 3183.7.

EXAMPLE 23Preparation of Compound 23

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Ala-NH₂ [SEQ.
ID. NO. 27]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 34% to 44% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 22.8 minutes. Electrospray Mass Spectrometry (M): calculated 3194.6; found 3197.6.

EXAMPLE 24Preparation of Compound 24

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
Pro Ser Ser Gly Ala Pro Pro Pro-NH₂ [SEQ. ID. NO. 28]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to

Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4099.6.

EXAMPLE 25

Preparation of Compound 25

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly
Pro Ser Ser Gly Ala Pro Pro Pro-NH₂ [SEQ. ID. NO. 29]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4042.5.

EXAMPLE 26

Preparation of Compound 26

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly

Pro Ser Ser Gly Ala Pro Pro-NH₂ [SEQ. ID. NO. 30]

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4002.4

EXAMPLE 27

Preparation of Compound 27

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly
Pro Ser Ser Gly Ala Pro Pro-NH₂ [SEQ. ID. NO. 31]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3945.4.

EXAMPLE 28Preparation of Compound 28

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly
Gly Pro Ser Ser Gly Ala Pro-NH₂ [SEQ. ID. NO. 32]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3905.3.

EXAMPLE 29Preparation of Compound 29

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly
Pro Ser Ser Gly Ala Pro-NH₂ [SEQ. ID. NO. 33]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved

from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3848.2.

EXAMPLE 30

Preparation of Compound 30

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 34]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3808.2.

EXAMPLE 31Preparation of Compound 31

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly
Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 35]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3751.1.

EXAMPLE 32Preparation of Compound 32

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
Pro Ser Ser Gly-NH₂ [SEQ. ID. NO. 36]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to

Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3737.1.

EXAMPLE 33

Preparation of Compound 33

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly
Pro Ser Ser Gly-NH₂ [SEQ. ID. NO. 37]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3680.1.

EXAMPLE 34

Preparation of Compound 34

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly

Pro Ser Ser-NH₂ [SEQ. ID. NO. 38]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3680.1

EXAMPLE 35

Preparation of Compound 35

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly
Pro Ser Ser-NH₂ [SEQ. ID. NO. 39]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3623.0.

EXAMPLE 36Preparation of Compound 36

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
Pro Ser-NH₂ [SEQ. ID. NO. 40]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3593.0

EXAMPLE 37Preparation of Compound 37

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly
Pro Ser-NH₂ [SEQ. ID. NO. 41]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved

from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3535.9

EXAMPLE 38

Preparation of Compound 38

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
Pro-NH₂ [SEQ. ID. NO. 42]

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3505.9.

EXAMPLE 39Preparation of Compound 39

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly
Pro-NH₂ [SEQ. ID. NO. 43]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3448.8.

EXAMPLE 40Preparation of Compound 40

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly-
NH₂ [SEQ. ID. NO. 44]

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to

Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3351.7.

EXAMPLE 41

Preparation of Compound 41

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly-NH₂
[SEQ. ID. NO. 45]

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3351.8.

EXAMPLE 42

Preparation of Compound 42

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu⁻
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly-NH₂

[SEQ. ID. NO. 46]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3294.7.

EXAMPLE 43

Preparation of Compound 43

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
tPro Ser Ser Gly Ala tPro tPro tPro-NH₂ [SEQ. ID. NO. 47]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Double couplings are required at residues 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass

Spectrometry (M): calculated 4197.1.

EXAMPLE 44

Preparation of Compound 44

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
Pro Ser Ser Gly Ala tPro tPro tPro-NH₂ [SEQ. ID. NO. 48]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Double couplings are required at residues 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4179.1.

EXAMPLE 45

Preparation of Compound 45

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
NMeala Ser Ser Gly Ala Pro Pro-NH₂ [SEQ. ID. NO. 49]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide

norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Double couplings are required at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3948.3.

EXAMPLE 46

Preparation of Compound 46

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
NMeala Ser Ser Gly Ala NMeala Nmeala-NH₂ [SEQ. ID. NO. 50]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Double couplings are required at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3840.1.

EXAMPLE 47Preparation of Compound 47

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
hPro Ser Ser Gly Ala hPro hPro-NH₂ [SEQ. ID. NO. 51]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Double couplings are required at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4050.1.

EXAMPLE 48Preparation of Compound 48

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
hPro Ser Ser Gly Ala hPro-NH₂ [SEQ. ID. NO. 52]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc⁻-protected amino acids (Applied Biosystems, Inc.), cleaved

from the resin, deprotected and purified in a similar way to Compound 1. A double coupling is required at residue 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3937.1

EXAMPLE 49

Preparation of Compound 49

Arg Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly
Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 53]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3827.2.

EXAMPLE 50Preparation of Compound 50

His Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly-
NH₂ [SEQ. ID. NO. 54]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3394.8.

EXAMPLE 51Preparation of Compound 51

His Gly Glu Gly Thr Naphthylala Thr Ser Asp Leu Ser Lys Gln
Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-
NH₂ [SEQ. ID. NO. 55]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to

Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3289.5.

EXAMPLE 52

Preparation of Compound 52

His Gly Glu Gly Thr Phe Ser Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂ [SEQ.
ID. NO. 56]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3280.7.

EXAMPLE 53

Preparation of Compound 53

His Gly Glu Gly Thr Phe Ser Thr Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂ [SEQ.

ID. NO. 57]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3294.7.

EXAMPLE 54

Preparation of Compound 54

His Gly Glu Gly Thr Phe Thr Ser Glu Leu Ser Lys Gln Met Ala
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂ [SEQ.
ID. NO. 58]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3250.7.

EXAMPLE 55Preparation of Compound 55

His Gly Glu Gly Thr Phe Thr Ser Asp pentylgly Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 59]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3253.5.

EXAMPLE 56Preparation of Compound 56

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Naphthylala Ile Glu Phe Leu Lys Asn-
NH₂ [SEQ. ID. NO. 60]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved

from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3289.5.

EXAMPLE 57

Preparation of Compound 57

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe tButylgly Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 61]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3183.4.

EXAMPLE 58Preparation of Compound 58

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Asp Phe Leu Lys Asn-NH₂ [SEQ.
ID. NO. 62]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3237.6.

EXAMPLE 59Preparation of Compound 59

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly
Pro Ser Ser-NH₂ [SEQ. ID. NO. 63]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to

Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3637.9.

EXAMPLE 60

Preparation of Compound 60

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly-NH₂
[SEQ. ID. NO. 64]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3309.7.

EXAMPLE 61

Preparation of Compound 61

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met Glu
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly

hPro Ser Ser Gly Ala hPro hPro-NH₂ [SEQ. ID. NO. 65]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Double couplings are required at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3711.1.

EXAMPLE 62

Preparation of Peptide having SEQ. ID. NO. 67

Compound 62, 4-imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys-NH⁺octanoyl Asn-NH₂ [SEQ. ID. NO. 67], is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 1. Fmoc-Lys-NH⁺octanoyl acid is used for coupling at position 27. Instead of using a protected amino acid for the final coupling at position 1, 4-imidazolylpropionic acid is coupled directly to the N-terminus of residues 2-28 on the resin. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in

Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3405.0

EXAMPLE 63

Preparation of Peptide having SEQ. ID. NO. 68

Compound 63, 4-imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys-NH^ooctanoyl Asn-NH₂ [SEQ. ID. NO. 68], is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 1. Fmoc-Lys-NH^ooctanoyl acid is used for coupling at position 27. Instead of using a protected amino acid for the final coupling at position 1, 4-imidazolylpropionic acid is coupled directly to the N-terminus of residues 2-28 on the resin. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3347.9

EXAMPLE 64

Preparation of Peptide having SEQ. ID. NO. 69

Compound 64, 4-imidazolylpropionyl-Gly Glu Gly Thr Phe

Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys-NH^toctanoyl Asn Gly Gly-NH₂ [SEQ. ID.
NO. 69], is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc
aminomethyl phenoxy acetamide norleucine MBHA resin
(Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids
(Applied Biosystems, Inc.), cleaved from the resin,
deprotected and purified in a similar way to Example 1.
Fmoc-Lys-NH^toctanoyl acid is used for coupling at position 27.
Instead of using a protected amino acid for the final
coupling at position 1, 4-imidazolylpropionic acid is coupled
directly to the N-terminus of residues 2-30 on the resin.
Used in analysis are Solvent A (0.1% TFA in water) and
Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30%
to 60% Solvent B in Solvent A over 30 minutes) of the
lyophilized peptide is then carried out to determine the
retention time of the product peptide. Electrospray Mass
Spectrometry (M): calculated 3519.0

EXAMPLE 65

Preparation of Peptide having SEQ. ID. NO. 70

Compound 65, 4-imidazolylpropionyl-Gly Glu Gly Thr Phe
Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys-NH^toctanoyl Asn Gly Gly-NH₂ [SEQ. ID.
NO. 70], is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc
aminomethyl phenoxy acetamide norleucine MBHA resin
(Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids
(Applied Biosystems, Inc.), cleaved from the resin,
deprotected and purified in a similar way to Example 1.
Fmoc-Lys-NH^toctanoyl acid is used for coupling at position 27.
Instead of using a protected amino acid for the final

coupling at position 1, 4-imidazolylpropionic acid is coupled directly to the N-terminus of residues 2-30 on the resin. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3451.9

EXAMPLE 66

Preparation of Peptide having SEQ. ID. NO. 71

Compound 66, 4-imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Asn Lys-NH⁸octanoyl-NH₂ [SEQ. ID. NO. 71], is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 1. Fmoc-Lys-NH⁸octanoyl acid is used for the initial coupling onto the resin at position 28. Instead of using a protected amino acid for the final coupling at position 1, 4-imidazolylpropionic acid is coupled directly to the N-terminus of protected residues 2-28 on the resin. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3405.0

EXAMPLE 67Preparation of Peptide having SEQ. ID. NO. 72

Compound 67, 4-imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Asn Lys-NH^toctanoyl-NH₂ [SEQ. ID. NO. 72], is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 1. Fmoc-Lys-NH^toctanoyl acid is used for the initial coupling onto the resin at position 28. Instead of using a protected amino acid for the final coupling at position 1, 4-imidazolylpropionic acid is coupled directly to the N-terminus of residues 2-28 on the resin. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3347.9

EXAMPLE 68Preparation of Peptide having SEQ. ID. NO. 73

Compound 68, 4-imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Asn Lys-NH^toctanoyl Gly Gly-NH₂ [SEQ. ID. NO. 73], is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin

(Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 1. Fmoc-Lys-NH^ooctanoyl acid is used for coupling at position 28. Instead of using a protected amino acid for the final coupling at position 1, 4-imidazolylpropionic acid is coupled directly to the N-terminus of protected residues 2-30 on the resin. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3519.0

EXAMPLE 69

Preparation of Peptide having SEQ. ID. NO. 74

Compound 69, 4-imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Asn Lys-NH^ooctanoyl Gly Gly-NH₂ [SEQ. ID. NO. 74], is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 1. Fmoc-Lys-NH^ooctanoyl acid is used for coupling at position 28. Instead of using a protected amino acid for the final coupling at position 1, 4-imidazolylpropionic acid is coupled directly to the N-terminus of residues 2-30 on the resin. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30%

to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3451.9

EXAMPLE 70

Preparation of C-terminal carboxylic acid peptides corresponding to the above C-terminal amide sequences for Compounds 1-3, 30-37, 40-42, 49-58 and 62-69

Compounds 1-23, 30-37, 40-42, 49-58 and 62-69 [SEQ. ID. NOS. 5-27, 34-41, 44-46, 53-62 and 67-74] are assembled on the so called Wang resin (p-alkoxybenzylalcohol resin (Bachem, 0.54 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

EXAMPLE 71

Preparation of C-terminal carboxylic acid peptides corresponding to the above C-terminal amide sequences for Compounds 24-29, 38, 39 and 43-48

Compounds 24-29, 38, 39, and 43-48 [SEQ. ID. NOS. 28-33, 42, 43 and 47-52] are assembled on the 2-chlorotritylchloride

resin (200-400 mesh), 2% DVB (Novabiochem, 0.4-1.0 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

EXAMPLES A TO E

Reagents Used

GLP-1(7-36)[NH₂] was purchased from Bachem (Torrance, CA). All other peptides were prepared using synthesis methods such as those described therein. All chemicals were of the highest commercial grade. The cAMP SPA immunoassay was purchased from Amersham. The radioligands were purchased from New England Nuclear (Boston, MA). RINm5f cells (American Type Tissue Collection, Rockville, MD) were grown in DME/F12 medium containing 10% fetal bovine serum and 2mM L-glutamine. Cells were grown at 37°C and 5% CO₂/95% humidified air and medium was replaced every 2 to 3 days. Cells were grown to confluence then harvested and homogenized using on a Polytron homogenizer. Cell homogenates were stored frozen at -70°C until used.

EXAMPLE A

GLP-1 Receptor Binding Studies

Receptor binding was assessed by measuring displacement of [¹²⁵I]GLP-1 or [¹²⁵I]exendin(9-39) from RINm5f membranes. Assay buffer contained 5 µg/ml bestatin, 1 µg/ml phosphoramidon, 1 mg/ml bovine serum albumin (fraction V), 1

mg/ml bacitracin, and 1 mM MgCl₂ in 20 mM HEPES, pH 7.4. To measure binding, 30 µg membrane protein (Bradford protein assay) was resuspended in 200 µl assay buffer and incubated with 60 pM [¹²⁵I]GLP-1 or [¹²⁵I]exendin(9-39) and unlabeled peptides for 120 minutes at 23°C in 96 well plates (Nagle Nunc, Rochester, NY). Incubations were terminated by rapid filtration with cold phosphatebuffered saline, pH 7.4, through polyethyleneimine-treated GF/B glass fiber filters (Wallac Inc., Gaithersburg, MD) using a Tomtec Mach II plate harvester (Wallac Inc., Gaithersburg, MD). Filters were dried, combined with scintillant, and radioactivity determined in a Betaplate liquid scintillant counter (Wallac Inc.).

Peptide samples were run in the assay as duplicate points at 6 dilutions over a concentration range of 10⁻⁶M to 10⁻¹²M to generate response curves. The biological activity of a sample is expressed as an IC₅₀ value, calculated from the raw data using an iterative curve-fitting program using a 4-parameter logistic equation (PrizmTM, GraphPAD Software). The results are shown in Table I.

TABLE I

<u>Compound</u>		<u>IC₅₀ (nM)</u>
Exendin-4	[SEQ. ID. NO. 2]	0.70
Compound 1	[SEQ. ID. NO. 5]	0.67
Compound 2	[SEQ. ID. NO. 6]	1.21
Compound 3	[SEQ. ID. NO. 7]	0.67
Compound 4	[SEQ. ID. NO. 8]	0.42
Compound 5	[SEQ. ID. NO. 9]	1.91
Compound 6	[SEQ. ID. NO. 10]	59.05
Compound 7	[SEQ. ID. NO. 11]	5.44
Compound 8	[SEQ. ID. NO. 12]	1.75
Compound 9	[SEQ. ID. NO. 13]	0.88
Compound 10	[SEQ. ID. NO. 14]	1.96
Compound 11	[SEQ. ID. NO. 15]	0.69
Compound 12	[SEQ. ID. NO. 16]	2.94
Compound 13	[SEQ. ID. NO. 17]	7.82
Compound 14	[SEQ. ID. NO. 18]	0.04
Compound 15	[SEQ. ID. NO. 19]	0.48
Compound 16	[SEQ. ID. NO. 20]	1.10
Compound 17	[SEQ. ID. NO. 21]	21.6
Compound 18	[SEQ. ID. NO. 22]	0.63
Compound 19	[SEQ. ID. NO. 23]	0.63
Compound 20	[SEQ. ID. NO. 24]	0.94
Compound 21	[SEQ. ID. NO. 25]	9.91
Compound 22	[SEQ. ID. NO. 26]	8.24
Compound 23	[SEQ. ID. NO. 27]	0.82

EXAMPLE BCyclase Activation Study

Assay buffer contained 10 μ M GTP, 0.75 mM ATP, 2.5 mM MgCl_2 , 0.5mM phosphocreatine, 12.5 U/ml creatine kinase, 0.4 mg/ml aprotinin, 1 μ M IBMX in 50 mM HEPES, pH 7.4. Membranes and peptides were combined in 100 μ l of assay buffer in 96 well filter-bottom plates (Millipore Corp., Bedford, MA). After 20 minutes incubation at 37°C, the assay was terminated by transfer of supernatant by filtration into a fresh 96 well plate using a Millipore vacuum manifold. Supernatant cAMP contents were quantitated by SPA immunoassay.

Peptide samples were run in the assay as triplicate points at 7 dilutions over a concentration range of 10^{-6} M to 10^{-12} M to generate response curves. The biological activity of a particular sample was expressed as an EC_{50} value, calculated as described above. Results are tabulated in Table II.

TABLE II

Compound	<u>EC₅₀ (nM)</u>
Exendin-4 [SEQ. ID. NO. 2]	0.23
Compound 1 [SEQ. ID. NO. 5]	0.3
Compound 2 [SEQ. ID. NO. 6]	0.79
Compound 3 [SEQ. ID. NO. 7]	2.35
Compound 4 [SEQ. ID. NO. 8]	0.22
Compound 5 [SEQ. ID. NO. 9]	9.85
Compound 6 [SEQ. ID. NO. 10]	79.4
Compound 7 [SEQ. ID. NO. 11]	63.6
Compound 8 [SEQ. ID. NO. 12]	6.8
Compound 9 [SEQ. ID. NO. 13]	1.68
Compound 10 [SEQ. ID. NO. 14]	5.37
Compound 11 [SEQ. ID. NO. 15]	0.48
Compound 12 [SEQ. ID. NO. 16]	15.55
Compound 13 [SEQ. ID. NO. 17]	79.6
Compound 14 [SEQ. ID. NO. 18]	1.11
Compound 15 [SEQ. ID. NO. 19]	1.05
Compound 16 [SEQ. ID. NO. 20]	5.12
Compound 17 [SEQ. ID. NO. 21]	43.6
Compound 18 [SEQ. ID. NO. 22]	0.76
Compound 19 [SEQ. ID. NO. 23]	3.68
Compound 20 [SEQ. ID. NO. 24]	5.25
Compound 21 [SEQ. ID. NO. 25]	45.1
Compound 22 [SEQ. ID. NO. 26]	20.43
Compound 23 [SEQ. ID. NO. 27]	3.05

EXAMPLE CDetermination of Blood Glucose Levels in db/db Mice

C57BLKS/J-m-db mice at least 3 months of age were utilized for the study. The mice were obtained from The Jackson Laboratory and allowed to acclimate for at least one week before use. Mice were housed in groups of ten at $22^{\circ} \pm 1^{\circ}\text{C}$ with a 12:12 light:dark cycle, with lights on at 6 a.m. All animals were deprived of food for 2 hours before taking baseline blood samples. Approximately 70 μl of blood was drawn from each mouse via eye puncture, after a light anesthesia with metophane. After collecting baseline blood samples, to measure plasma glucose concentrations, all animals receive subcutaneous injections of either vehicle (10.9% NaCl), exendin-4 or test compound (1 μg) vehicle. Blood samples were drawn again, using the same procedure, after exactly one hour from the injections, and plasma glucose concentrations were measured.

For each animal, the % change in plasma value, from baseline value, was calculated. The percent decrease in plasma glucose after one hour is shown in Table III.

TABLE III

<u>Test Compound</u>	<u>% drop in glucose</u>	
Exendin-4 [SEQ. ID. NO. 2]	39%	(n = 78)
Compound 2 [SEQ. ID. NO. 6]	38%	(n = 4)
Compound 3 [SEQ. ID. NO. 7]	49%	(n = 4)
Compound 4 [SEQ. ID. NO. 8]	27%	(n = 4)
Compound 5 [SEQ. ID. NO. 9]	47%	(n = 4)
Compound 6 [SEQ. ID. NO. 10]	40%	(n = 5)
Compound 7 [SEQ. ID. NO. 11]	31%	(n = 4)
Compound 8 [SEQ. ID. NO. 12]	44%	(n = 4)
Compound 9 [SEQ. ID. NO. 13]	41%	(n = 4)
Compound 10 [SEQ. ID. NO. 14]	46%	(n = 4)
Compound 11 [SEQ. ID. NO. 15]	40%	(n = 4)
Compound 12 [SEQ. ID. NO. 16]	53%	(n = 4)
Compound 13 [SEQ. ID. NO. 17]	45%	(n = 4)
Compound 14 [SEQ. ID. NO. 18]	54%	(n = 4)
Compound 15 [SEQ. ID. NO. 19]	45%	(n = 4)
Compound 16 [SEQ. ID. NO. 20]	54%	(n = 4)
Compound 17 [SEQ. ID. NO. 21]	45%	(n = 4)
Compound 18 [SEQ. ID. NO. 22]	50%	(n = 4)
Compound 19 [SEQ. ID. NO. 23]	48%	(n = 4)
Compound 20 [SEQ. ID. NO. 24]	37%	(n = 4)
Compound 21 [SEQ. ID. NO. 25]	30%	(n = 4)
Compound 22 [SEQ. ID. NO. 26]	46%	(n = 4)
Compound 23 [SEQ. ID. NO. 27]	42%	(n = 4)

EXAMPLE DDose Response Determination of Blood Glucose Levels in
db/db Mice

C57BLKS/J-m-db/db mice, at least 3 months of age were utilized for the study. The mice were obtained from The Jackson Laboratory and allowed to acclimate for at least one week before use. Mice were housed in groups of ten at 22°C 1°C with a 12:12 light:dark cycle, with lights on at 6 a.m.

All animals were deprived of food for 2 hours before taking baseline blood samples. Approximately 70 µl of blood was drawn from each mouse via eye puncture, after a light anesthesia with metophane. After collecting baseline blood samples, to measure plasma glucose concentrations, all animals receive subcutaneous injections of either vehicle, exendin-4 or test compound in concentrations indicated. Blood samples were drawn again, using the same procedure, after exactly one hour from the injections, and plasma glucose concentrations were measured.

For each animal, the % change in plasma value, from baseline value, was calculated and a dose dependent relationship was evaluated using Graphpad Prism™ software.

Figure 5 depicts the effects of varying doses of exendin-4 [SEQ. ID. NO. 2] and Compound 3 [SEQ. ID. NO. 7] on plasma glucose levels. Exendin-4 had an ED₅₀ of 0.01 µg per mouse and Compound 3 had an ED₅₀ of 0.04 µg per mouse.

EXAMPLE EGastric Emptying

The following study was carried out to examine the effects of exendin-4 and exendin agonist compounds of the present invention on gastric emptying in rats. This experiment followed a modification of the method of

Scarpignato, et al., Arch. Int. Pharmacodyn. Ther.
246:286-94, 1980.

Male Harlan Sprague Dawley (HSD) rats were used. All animals were housed at 22.7 ± 0.8 C in a 12:12 hour light:dark cycle (experiments being performed during the light cycle) and were fed and watered *ad libitum* (Diet LM-485, Teklad, Madison, WI). Exendin-4 was synthesized according to standard peptide synthesis methods. The preparation of Compounds 2, 3, 10 and 13 [SEQ. ID. NOS. 6, 7, 14 and 17] is described in Examples 2, 3, 10 and 13, respectively.

The determination of gastric emptying by the method described below was performed after a fast of ~20 hours to ensure that the stomach contained no chyme that would interfere with spectrophotometric absorbance measurements.

Conscious rats received by gavage, 1.5ml of an acaloric gel containing 1.5% methyl cellulose (M-0262, Sigma Chemical Co, St Louis, MO) and 0.05% phenol red indicator. Twenty minutes after gavage, rats were anesthetized using 5% halothane, the stomach exposed and clamped at the pyloric and lower esophageal sphincters using artery forceps, removed and opened into an alkaline solution which was made up to a fixed volume. Stomach content was derived from the intensity of the phenol red in the alkaline solution, measured by absorbance at a wavelength of 560 nm. In separate experiments on 7 rats, the stomach and small intestine were both excised and opened into an alkaline solution. The quantity of phenol red that could be recovered from the upper gastrointestinal tract within 20 minutes of gavage was $89 \pm 4\%$; dye which appeared to bind irrecoverably to the gut luminal surface may have accounted for the balance. To account for a maximal dye recovery of less than 100%, percent of stomach contents remaining after 20 min were expressed as a fraction

of the gastric contents recovered from control rats sacrificed immediately after gavage in the same experiment. Percent gastric contents remaining = (absorbance at 20 min)/(absorbance at 0 min) x 100.

In baseline studies, with no drug treatment, gastric emptying over 20 min was determined. In dose-response studies, rats were treated with 0.01, 0.1, 0.3, 1, 10 and 100 µg of exendin-4, and 0.1, 0.3, 1, 10 and 100 µg of Compounds 2, 3, 10 and 13 [SEQ. ID. NOS. 6, 7, 14 and 17].

The results, shown in Figures 6-9, show that the exendin agonists, Compounds 2, 3, 10 and 13, are potent inhibitors of gastric emptying. The EC₅₀ for exendin-4 was 0.27 µg. The EC₅₀ for Compound 2 was 0.79 µg, for Compound 3 was 0.34 µg, for Compound 10 was 0.27 µg, and for Compound 13 was 60 µg.

We claim:

1. A peptide compound of the formula (I) [SEQ. ID. NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁; wherein

Xaa₁ is His, Arg or Tyr;
Xaa₂ is Ser, Gly, Ala or Thr;
Xaa₃ is Asp or Glu;
Xaa₅ is Ala or Thr;
Xaa₆ is Ala, Phe, Tyr or naphthylalanine;
Xaa₇ is Thr or Ser;
Xaa₈ is Ala, Ser or Thr;
Xaa₉ is Asp or Glu;
Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
Xaa₁₁ is Ala or Ser;
Xaa₁₂ is Ala or Lys;
Xaa₁₃ is Ala or Gln;
Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
Xaa₁₅ is Ala or Glu;
Xaa₁₆ is Ala or Glu;
Xaa₁₇ is Ala or Glu;
Xaa₁₉ is Ala or Val;
Xaa₂₀ is Ala or Arg;
Xaa₂₁ is Ala or Leu;
Xaa₂₂ is Phe, Tyr or naphthylalanine;
Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met;

Xaa₂₄ is Ala, Glu or Asp;

Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;

Xaa₂₆ is Ala or Leu;

Xaa₂₇ is Ala or Lys;

Xaa₂₈ is Ala or Asn;

Z₁ is -OH,

-NH₂,

Gly-Z₂,

Gly Gly -Z₂

Gly Gly Xaa₃₁-Z₂,

Gly Gly Xaa₃₁ Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂;

wherein

Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently

selected from the group consisting of Pro,

homoproline, 3Hyp, 4Hyp, thioproline,

N-alkylglycine, N-alkylpentylglycine and

N-alkylalanine; and

Z₂ is -OH or -NH₂;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈,

Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀,

Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇, and Xaa₂₈ are Ala; and

pharmaceutically acceptable salts thereof.

2. A compound according to claim 1 wherein Xaa₁ is His or Tyr.

3. A compound according to claim 2 wherein Xaa₁ is His.
4. A compound according to claim 2 wherein Xaa₂ is Gly.
5. A compound according to claim 4 wherein Xaa₁₄ is Leu, pentylglycine or Met.
6. A compound according to claim 5 wherein Xaa₂₅ is Trp or Phe.
7. A compound according to claim 6 wherein Xaa₆ is Phe or naphthylalanine; and Xaa₂₂ is Phe or naphthylalanine; Xaa₂₃ is Ile or Val.
8. A compound according to claim 7 wherein Z₁ is -NH₂.
9. A compound according to claim 7 wherein Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected from the group consisting of Pro, homoproline, thioproline and N-alkylalanine.
10. A compound according to claim 9 wherein Z₂ is -NH₂.
11. A compound according to claim 1 wherein Xaa₂ is Gly.
12. A compound according to claim 1 wherein Xaa₁₄ is Leu, pentylglycine or Met.
13. A compound according to claim 1 wherein Xaa₂₅ is Trp or Phe.
14. A compound according to claim 1 wherein Xaa₆ is Ph⁻ or naphthylalanine; Xaa₂₂ is Phe or naphthylalanine; Xaa₂₃ is

Ile or Val.

15. A compound according to claim 1 wherein Z_1 is $-NH_2$.

16. A compound according to claim 1 wherein Xaa_{31} , Xaa_{36} , Xaa_{37} and Xaa_{38} are independently selected from the group consisting of Pro, homoproline, thioproline and N-alkylalanine.

17. A compound according to claim 1 wherein Z_2 is $-NH_2$.

18. A compound according to claim 1 which has an amino acid sequence selected from SEQ. ID. NOS. 5 to 65.

19. A peptide compound of the formula (I) [SEQ. ID. NO. 4]:

Xaa_1 Xaa_2 Xaa_3 Gly Xaa_5 Xaa_6 Xaa_7 Xaa_8 Xaa_9 Xaa_{10}
 Xaa_{11} Xaa_{12} Xaa_{13} Xaa_{14} Xaa_{15} Xaa_{16} Xaa_{17} Ala Xaa_{18} Xaa_{19}
 Xaa_{20} Xaa_{21} Xaa_{22} Xaa_{23} Xaa_{24} Xaa_{25} Xaa_{26} Xaa_{27} $Xaa_{28}-Z_1$;

wherein

Xaa_1 is His or Arg;

Xaa_2 is Gly or Ala;

Xaa_3 is Asp or Glu;

Xaa_5 is Ala or Thr;

Xaa_6 is Ala, Phe or naphthylalanine;

Xaa_7 is Thr or Ser;

Xaa_8 is Ala, Ser or Thr;

Xaa_9 is Asp or Glu;

Xaa_{10} is Ala, Leu or pentylglycine;

Xaa_{11} is Ala or Ser;

Xaa₁₂ is Ala or Lys;
Xaa₁₃ is Ala or Gln;
Xaa₁₄ is Ala, Leu or pentylglycine;
Xaa₁₅ is Ala or Glu;
Xaa₁₆ is Ala or Glu;
Xaa₁₇ is Ala or Glu;
Xaa₁₉ is Ala or Val;
Xaa₂₀ is Ala or Arg;
Xaa₂₁ is Ala or Leu;
Xaa₂₂ is Phe or naphthylalanine;
Xaa₂₃ is Ile, Val or tert-butylglycine;
Xaa₂₄ is Ala, Glu or Asp;
Xaa₂₅ is Ala, Trp, or Phe;
Xaa₂₆ is Ala or Leu;
Xaa₂₇ is Ala or Lys;
Xaa₂₈ is Ala or Asn;
Z₁ is -OH,
-NH₂,
Gly-Z₂,
Gly Gly -Z₂,
Gly Gly Xaa₃₁-Z₂,
Gly Gly Xaa₃₁ Ser-Z₂,
Gly Gly Xaa₃₁ Ser Ser-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or Gly Gly
Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂;
Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected
from the group consisting of Pro, homoproline,
thioprolin and N-methylalanine; and
Z₂ is -OH or -NH₂;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈ are Ala; and pharmaceutically acceptable salts thereof.

20. A compound according to claim 19 which has an amino acid sequence selected from SEQ. ID. NOS. 6-19.

21. A composition comprising a compound of any of claims 1 to 19 in a pharmaceutically acceptable carrier.

22. A composition comprising a compound of claim 20 in a pharmaceutically acceptable carrier.

23. A method for the treatment of diabetes mellitus comprising the administration of a therapeutically effective amount of a compound according to claim 1.

24. A method for the treatment of diabetes mellitus comprising the administration of a therapeutically effective amount of a compound according to claim 18.

25. A method for the treatment of diabetes mellitus comprising the administration of a therapeutically effective amount of a compound according to claim 19.

26. A method for the treatment of diabetes mellitus comprising the administration of a therapeutically effective amount of a compound according to claim 20.

27. The method of claim 23 further comprising the administration of a therapeutically effective amount of an insulin.

28. The method of claim 24 further comprising the administration of a therapeutically effective amount of an insulin.

29. The method of claim 25 further comprising the administration of a therapeutically effective amount of an insulin.

30. The method of claim 26 further comprising the administration of a therapeutically effective amount of an insulin.

31. A method for the treatment of a hyperglycemic condition in a mammal comprising the step of administering a therapeutically effective amount of a compound according to claim 1.

32. A method for the treatment of a hyperglycemic condition in a mammal comprising the step of administering a therapeutically effective amount of a compound according to claim 18.

33. A method for the treatment of a hypoglycemic condition in a mammal comprising the step of administering a therapeutically effective amount of a compound according to claim 19.

34. A method for the treatment of a hypoglycemic condition in a mammal comprising the step of administering a therapeutically effective amount of a compound according to claim 20.

35. A peptide compound of the formula (II) [SEQ. ID. NO. 66]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ X_{1-Z}; wherein

Xaa₁ is His, Arg or Tyr or 4-imidazopropionyl;

Xaa, is Ser, Gly, Ala or Thr;

Xaa₃ is Asp or Glu;

Xaa₅ is Ala or Thr;

Xaa₆ is Ala, Phe, Tyr or naphthylalanine;

Xaa, is Thr or Ser;

Xaa₈ is Ala, Ser or Thr;

Xaa₉ is Asp or Glu;

Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;

Xaa₁₁ is Ala or Ser;

Xaa₁₂ is Ala or Lys;

Xaa₁₃ is Ala or Gln;

Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;

Xaa₁₅ is Ala or Glu;

Xaa₁₆ is Ala or Glu;

Xaa₁₇ is Ala or Glu;

Xaa₁₉ is Ala or Val;

Xaa₂₀ is Ala or Arg;

Xaa₂₁ is Ala, Leu or Lys-NH^e-R where R is Lys, Arg, C₁-C₁₀ straight chain or branched alkanoyl or cycloalkylkanoyl;

Xaa₂₂ is Phe, Tyr or naphthylalanine;

Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met;

Xaa₂ is Ala, Glu or Asp;

Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;

Xaa₂₆ is Ala or Leu;

X₁ is Lys Asn, Asn Lys, Lys-NH^e-R Asn, Asn Lys-NH^e-R, Lys-NH^e-R Ala, Ala Lys-NH^e-R where R is Lys, Arg, C₁-C₁₀ straight chain or branched alkanoyl or cycloalkylalkanoyl

Z₁ is -OH,

-NH₂,

Gly-Z₂,

Gly Gly-Z₂

Gly Gly Xaa₃₁-Z₂,

Gly Gly Xaa₃₁ Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂; wherein

Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently

selected from the group consisting of Pro,

homoproline, 3Hyp, 4Hyp, thioproline,

N-alkylglycine, N-alkylpentylglycine and

N-alkylalanine; and

Z₂ is -OH or -NH₂;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, and Xaa₂₆ are Ala; and pharmaceutically acceptable salts thereof.

36. A compound according to claim 35 wherein Xaa₁ is His, Tyr or 4-imidazopropionyl.

37. A compound according to claim 36 wherein Xaa₁ is His.

38. A compound according to claim 36 wherein Xaa₁ is 4-imidazopropionyl.

39. A compound according to claim 35 wherein Xaa₂ is Gly.

40. A compound according to claim 35 wherein Xaa₁₄ is Leu, pentylglycine or Met.

41. A compound according to claim 35 wherein Xaa₂₅ is Trp or Phe.

42. A compound according to claim 35 wherein Xaa₆ is Phe or naphthylalanine; Xaa₂₂ is Phe or naphthylalanine; and Xaa₂₃ is Ile or Val.

43. A compound according to claim 35 wherein Z₁ is -NH₂.

44. A compound according to claim 35 wherein Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently selected from the group consisting of Pro, homoproline, thioproline and N-alkylalanine.

45. A compound according to claim 35 wherein Z₂ is -NH₂.

46. A compound according to claim 35 wherein X₁ is Lys Asn, Lys-NH^ε-R Asn, or Lys-NH^ε-R Ala where R is Lys, Arg, C₁-C₁₀ straight chain or branched alkanoyl.

47. A compound according to claim 35 wherein Xaa₂₁ is Lys-NH^ε-R where R is Lys, Arg, C₁-C₁₀ straight chain or branched alkanoyl or cycloalkylkanoyl.

48. A compound according to claim 35 wherein said compound has an amino acid sequence selected from SEQ. ID. NOS. 67-74.

49. A composition comprising a compound of any of claims 35-47 in a pharmaceutically acceptable carrier.

50. A composition comprising a compound of claim 48 in a pharmaceutically acceptable carrier.

01/09

EXENDIN-3

His	Ser	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu
1				5					10					15	
Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser
			20					25					30		
Ser	Gly	Ala	Pro	Pro	Pro	Ser-NH ₂									
			35												

Fig. 1

EXENDIN-4

His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu
				5					10					15	
Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser
			20					25					30		
Ser	Gly	Ala	Pro	Pro	Pro	Ser-NH ₂									
			35												

*Fig. 2*GLP-1 (GLP-1[7-36]NH₂)

His	Ala	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Val	Ser	Ser	Tyr	Leu	Glu	Gly
				5					10					15	
Gln	Ala	Ala	Lys	Glu	Phe	Ile	Ala	Trp	Leu	Val	Lys	Gly	Arg-NH ₂		
			20					25					30		

Fig. 3

SUBSTITUTE SHEET (RULE 26)

Anticore Acid Position			1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38					
Compound 1	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	NH2															
Compound 2	His	Gly	Gly	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH2																	
Compound 3	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 4	His	Ala	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 5	His	Gly	Glu	Gly	Ala	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 6	His	Gly	Glu	Gly	Thr	Ala	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 7	His	Gly	Glu	Gly	Thr	Phe	Thr	Ala	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 8	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 9	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ala	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 10	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 11	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Ala	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																
Compound 12	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Ala	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																
Compound 13	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 14	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 15	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 16	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 17	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 18	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Ala	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 19	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 20	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Ala	Leu	Lys	Asn	NH2																	
Compound 21	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Ala	Lys	Asn	NH2																	
Compound 22	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 23	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2																	
Compound 24	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	Pro	Pro	NH2					
Compound 25	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	Pro	Pro	Pro	NH2				
Compound 26	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	Pro	Pro	NH2					
Compound 27	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	Pro	Pro	NH2					
Compound 28	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	Pro	NH2						
Compound 29	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	NH2							
Compound 30	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	NH2							
Compound 31	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	NH2							
Compound 32	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	NH2							
Compound 33	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	NH2							
Compound 34	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	NH2							
Compound 35	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Ser	Gly	Ala	Pro	Ala	NH2							

Fig. 4A

03/09


Amino Acid Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	
Compound 36 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	NH2						
Compound 37 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	NH2						
Compound 38 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	NH2							
Compound 39 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	NH2							
Compound 40 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	NH2								
Compound 41 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	NH2									
Compound 42 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	NH2									
Compound 43 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	hPro	Ser	Ser	Gly	Ala	hPro	hPro	NH2	
Compound 44 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	hPro	hPro	NH2	
Compound 45 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Nme	Ser	Ser	Gly	Ala	Pro	Pro	NH2	
Compound 46 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Nme	Ser	Ser	Gly	Ala	Nme	Nme	NH2	
Compound 47 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	hPro	Ser	Ser	Gly	Ala	hPro	hPro	NH2	
Compound 48 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	hPro	Ser	Ser	Gly	Ala	hPro	NH2		
Compound 49 Arg	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	NH2			
Compound 50 His	Gly	Asp	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	NH2								
Compound 51 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2										
Compound 52 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH2										
Compound 53 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH2										
Compound 54 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Glu	Leu	Ser	Lys	Gln	Met	Ala	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH2										
Compound 55 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Gly	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2										
Compound 56 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg	Leu	naph	Ile	Glu	Phe	Leu	Lys	Asn	NH2										
Compound 57 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH2										
Compound 58 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	As	Phe	Leu	Lys	Asn	NH2										
Compound 59 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	NH2					
Compound 60 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	NH2									
Compound 61 His	Gly	Glu	Gly	Gly	Thr	Thr	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	hPro	Ser	Ser	Gly	Ala	hPro	hPro	NH2	

Fig. 4B

SUBSTITUTE SHEET (RULE 26)

04/09

Glucose lowering effect in db/db mice at
1 hr time point

*Fig. 5*

SUBSTITUTE SHEET (RULE 26)

05/09




Fig. 6

06/09




Fig. 7

07/09




Fig. 8

SUBSTITUTE SHEET (RULE 26)

BEST AVAILABLE COPY

08/09




Fig. 9

Compound
No.

- 62 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys-NH⁺octanoyl Asn-NH₂ [SEQ. ID NO. 67]
- 63 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Lys-NH⁺octanoyl Asn-NH₂ [SEQ. ID NO. 68]
- 64 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Lys-NH⁺octanoyl Asn Gly Gly-NH₂ [SEQ. ID NO. 69]
- 65 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val
Arg Leu Phe Ile Glu Phe Leu Lys-NH⁺octanoyl Asn Gly Gly-NH₂ [SEQ. ID NO. 70]
- 66 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Asn Lys-NH⁺octanoyl-NH₂ [SEQ. ID NO. 71]
- 67 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Asn Lys-NH⁺octanoyl-NH₂ [SEQ. ID NO. 72]
- 68 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Trp Leu Asn Lys-NH⁺octanoyl Gly Gly-NH₂ [SEQ. ID NO. 73]
- 69 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu
Phe Ile Glu Phe Leu Asn Lys-NH⁺octanoyl Gly Gly-NH₂ [SEQ. ID NO. 74]

Fig. 10